

# Summary

The summary file contains summary information for all the raw files processed with a single MaxQuant run. The summary information consists of some MaxQuant parameters, information of the raw file contents, and statistics on the peak detection. Based on this file a quick overview can be gathered on the quality of the data in the raw file.

The last row in this file contains the summary information for each column on each of the processed files.

Name	Separator	Description
Raw file		The raw file processed.
Experiment		Experiment name assigned to this LC-MS run in the experimental design.
Fraction		Fraction assigned to this LC-MS run in the experimental design.
Enzyme		The protease used to digest the protein sample.
Enzyme mode		The protease used to digest the protein sample.
Enzyme first search		The protease used for the first search.
Enzyme mode first search		The protease used for the first search.
Use enzyme first search		Marked with '+' when a different protease setup was used for the first search.
Variable modifications		The variable modification(s) used during the identification of peptides.
Fixed modifications		The fixed modification(s) used during the identification of peptides.
Multi modifications		The multi modification(s) used during the identification of peptides.
Variable modifications first search		The variable modification(s) used during the first search.
Use variable modifications first search		Marked with '+' when different variable modifications were used for the first search.
Requantify		The number of labels used.
Multiplicity		The number of labels used.
Max. missed cleavages		The maximum allowed number of missed cleavages.
Max. labeled AAs		The maximum allowed of labeled amino acids in a peptide amino acid sequence.
Labels0		The labels used in the labeling experiment. Allowed values for X: 0=light; 1=medium; 2=heavy label partner.
Labels1		The labels used in the labeling experiment. Allowed values for X: 0=light; 1=medium; 2=heavy label partner.
LC-MS run type		The type of LC-MS run. Usually it will be 'Standard' which refers to a conventional shotgun proteomics run with data-dependent MS/MS.
Time-dependent recalibration		When marked with '+', time-dependent recalibration was applied to improve the data quality.
MS		The number of MS spectra recorded in this raw file.
MS/MS		The number of MS/MS spectra recorded in this raw file.
MS3		The number of MS3 spectra recorded in this raw file.
MS/MS submitted		The number of tandem MS spectra submitted for analysis.
MS/MS submitted (SIL)		The number of tandem MS spectra submitted for analysis, where the precursor ion was detected as part of a labeling cluster.
MS/MS submitted (ISO)		The number of tandem MS spectra submitted for analysis, where the precursor ion was detected as an isotopic pattern.
MS/MS submitted (PEAK)		The number of tandem MS spectra submitted for analysis, where the precursor ion was detected as a single peak.
MS/MS identified		The total number of identified tandem MS spectra.
MS/MS identified (SIL)		The total number of identified tandem MS spectra, where the precursor ion was detected as part of a labeling cluster.
MS/MS identified (ISO)		The total number of identified tandem MS spectra, where the precursor ion was detected as an isotopic pattern.
MS/MS identified (PEAK)		The total number of identified tandem MS spectra, where the precursor ion was detected as a single peak.
MS/MS identified [%]		The percentage of identified tandem MS spectra.
MS/MS identified (SIL) [%]		The percentage of identified tandem MS spectra, where the precursor ion was detected as part of a labeling cluster.
MS/MS identified (ISO) [%]		The percentage of identified tandem MS spectra, where the precursor ion was detected as an isotopic pattern.

MS/MS identified (PEAK) [%]		The percentage of identified tandem MS spectra, where the precursor ion was detected as a single peak.
Peptide sequences identified		The total number of unique peptide amino acid sequences identified from the recorded tandem mass spectra.
Peaks		The total number of peaks detected in the full scans.
Peaks sequenced		The total number of peaks sequenced by tandem MS.
Peaks sequenced [%]		The percentage of peaks sequenced by tandem MS.
Peaks repeatedly sequenced		The total number of peaks repeatedly sequenced (i.e. 1 or more times) by tandem MS.
Peaks repeatedly sequenced [%]		The percentage of peaks repeatedly sequenced (i.e. 1 or more times) by tandem MS.
Isotope patterns		The total number of detected isotope patterns.
Isotope patterns sequenced		The total number of isotope patterns sequenced by tandem MS.
Isotope patterns sequenced (z>1)		The total number of isotope patterns sequenced by tandem MS with a charge state of 2 or more.
Isotope patterns sequenced [%]		The percentage of isotope patterns sequenced by tandem MS.
Isotope patterns sequenced (z>1) [%]		The percentage of isotope patterns sequenced by tandem MS with a charge state of 2 or more.
Isotope patterns repeatedly sequenced		The total number of isotope patterns repeatedly sequenced (i.e. 1 or more times) by tandem MS.
Isotope patterns repeatedly sequenced [%]		The percentage of isotope patterns repeatedly sequenced (i.e. 1 or more times) by tandem MS.
Multiplets		The total number of detected labeling pairs.
Multiplets z=1		The total number of detected labeling pairs with a charge state of 1.
Multiplets z=2		The total number of detected labeling pairs with a charge state of 2.
Multiplets z=3		The total number of detected labeling pairs with a charge state of 3.
Multiplets z=4		The total number of detected labeling pairs with a charge state of 4.
Multiplets z=5		The total number of detected labeling pairs with a charge state of 5.
Multiplets z=6		The total number of detected labeling pairs with a charge state of 6.
Multiplets z=7		The total number of detected labeling pairs with a charge state of 7.
Multiplets sequenced		The total number of labeling pairs sequenced by tandem MS.
Multiplets sequenced [%]		The percentage of labeling pairs sequenced by tandem MS.
Multiplets repeatedly sequenced		The total number of labeling pairs repeatedly sequenced (i.e. 1 or more times) by tandem MS.
Multiplets repeatedly sequenced [%]		The percentage of labeling pairs repeatedly sequenced (i.e. 1 or more times) by tandem MS.
Multiplets identified		The total number of labeling pairs identified.
Multiplets identified [%]		The percentage of labeling pairs identified.
Recalibrated		When marked with '+', the masses taken from the raw file were recalibrated.
Av. absolute mass deviation [ppm]		The average absolute mass deviation found comparing to the identification mass in parts per million.
Mass standard deviation [ppm]		The standard deviation of the mass deviation found comparing to the identification mass in parts per million.
Av. absolute mass deviation [mDa]		The average absolute mass deviation found comparing to the identification mass in milli-Dalton.
Mass standard deviation [mDa]		The standard deviation of the mass deviation found comparing to the identification mass in milli-Dalton.

# Evidence

The evidence file combines all the information about the identified peptides and normally is the only file required for processing the results. Additional information about the peptides, modifications, proteins, etc. can be found in the other files by unique identifier linkage.

Name	Separator	Description
Sequence		The identified AA sequence of the peptide.
Length		The length of the sequence stored in the column 'Sequence'.
K Count		The number of instances of K contained within the sequence. The value for this can reliably be determined in the case of labeling partners based on the distance between the partners. These counts are used to solidify the peptide identification process.
Modifications		Post-translational modifications contained within the identified peptide sequence.
Modified sequence		Sequence representation including the post-translational modifications (abbreviation of the modification in brackets before the modified AA). The sequence is always surrounded by underscore characters ('_').
Acetyl (K) Probabilities		Sequence representation of the peptide including PTM positioning probabilities ([0..1], where 1 is best match) for 'Acetyl (K)'.
Oxidation (M) Probabilities		Sequence representation of the peptide including PTM positioning probabilities ([0..1], where 1 is best match) for 'Oxidation (M)'.
Acetyl (K) Score Diffs		Sequence representation for each of the possible PTM positions in each possible configuration, the difference is calculated between the identification score with the PTM added to that position and the best scoring identification where no PTM is added to that position. When this value is negative, it is unlikely that the particular modification is located at this position.
Oxidation (M) Score Diffs		Sequence representation for each of the possible PTM positions in each possible configuration, the difference is calculated between the identification score with the PTM added to that position and the best scoring identification where no PTM is added to that position. When this value is negative, it is unlikely that the particular modification is located at this position.
Acetyl (K)		The number of occurrences of the modification 'Acetyl (K)'.
Acetyl (Protein N-term)		The number of occurrences of the modification 'Acetyl (Protein N-term)'.
Oxidation (M)		The number of occurrences of the modification 'Oxidation (M)'.
Missed cleavages		Number of missed enzymatic cleavages.
Proteins		The identifiers of the proteins this particular peptide is associated with.
Leading proteins		The identifiers of the proteins in the proteinGroups file, with this protein as best match, this particular peptide is associated with. When multiple matches are found here, the best scoring protein can be found in the 'Leading Razor Protein' column.
Leading razor protein		The identifier of the best scoring protein, from the proteinGroups file this, this peptide is associated to.
Gene names		Names of genes this peptide is associated with.
Protein names		Names of proteins this peptide is associated with.
Type		The type of the feature. 'MSMS' for an MS/MS spectrum without an MS1 isotope pattern assigned. 'ISO-MSMS' MS1 isotope cluster identified by MS/MS. 'MULTI-MSMS' MS1 labeling cluster identified by MS/MS. 'MULTI-SECPEP' MS1 labeling cluster identified by MS/MS as second peptide. 'MULTI-MATCH' MS1 labeling cluster identified by matching between runs. In case of label-free data there is no difference between 'MULTI' and 'ISO'.
Labeling State		Labeling state of the precursor isotope pattern used to identify the peptide.
Raw file		The name of the RAW-file the mass spectral data was derived from.
Fraction		The fraction in which this peptide was detected.
Experiment		
MS/MS m/z		The m/z used for fragmentation (not necessarily the mono-isotopic m/z).
Charge		The charge-state of the precursor ion.
m/z		The recalibrated mass-over-charge value of the precursor ion.

Mass		The predicted monoisotopic mass of the identified peptide sequence.
Uncalibrated - Calibrated m/z [ppm]		The difference between the uncalibrated and recalibrated mass-over-charge value of the precursor ion measured in parts-per-million. This gives an indication of the mass drift in the original data, which was automatically corrected by MaxQuant.
Uncalibrated - Calibrated m/z [Da]		The difference between the uncalibrated and recalibrated mass-over-charge value of the precursor ion measured in parts-per-million. This gives an indication of the mass drift in the original data, which was automatically corrected by MaxQuant.
Mass error [ppm]		Mass error of the recalibrated mass-over-charge value of the precursor ion in comparison to the predicted monoisotopic mass of the identified peptide sequence in parts per million.
Mass error [Da]		Mass error of the recalibrated mass-over-charge value of the precursor ion in comparison to the predicted monoisotopic mass of the identified peptide sequence in milli-Dalton.
Uncalibrated mass error [ppm]		Mass error of the uncalibrated mass-over-charge value of the precursor ion in comparison to the predicted monoisotopic mass of the identified peptide sequence.  Note: This column can contain missing values (denoted as NaN).
Uncalibrated mass error [Da]		Mass error of the uncalibrated mass-over-charge value of the precursor ion in comparison to the predicted monoisotopic mass of the identified peptide sequence.  Note: This column can contain missing values (denoted as NaN).
Max intensity m/z 0		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Max intensity m/z 1		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Retention time		The uncalibrated retention time in minutes in the elution profile of the precursor ion.
Retention length		The total retention time length of the peak (last time point - first time point).
Calibrated retention time		The recalibrated retention time in minutes in the elution profile of the precursor ion.
Calibrated retention time start		The recalibrated retention time start in minutes in the elution profile of the precursor ion.
Calibrated retention time finish		The recalibrated retention finish in minutes in the elution profile of the precursor ion.
Retention time calibration		The difference in minutes between the uncalibrated and recalibrated retention time. This gives an indication of the retention time drift in the original data, which was automatically corrected by MaxQuant.  Note: This column can contain missing values (NaN).
Match time difference		When the option match between runs is used in MaxQuant, this value indicates the time difference between the feature from the raw file it was taken from and the feature from the raw file it was matched to.
Match m/z difference		When the option match between runs is used in MaxQuant, this value indicates the m/z difference between the feature from the raw file it was taken from and the feature from the raw file it was matched to.
Match q-value		This is the q-value for features that have been identified by 'matching between runs'.
Match score		The andromeda score of the MS/MS identification that is the source of this identification by 'matching between runs'.
Number of data points		The number of data points (peak centroids) collected for this peptide feature.
Number of scans		The number of MS scans that the 3d peaks of this peptide feature are overlapping with.
Number of isotopic peaks		The number of isotopic peaks contained in this peptide feature.
PIF		Short for Parent Ion Fraction; indicates the fraction the target peak makes up of the total intensity in the inclusion window.
Fraction of total spectrum		The percentage the ion intensity makes up of the total intensity of the whole spectrum.
Base peak fraction		The percentage the parent ion intensity in comparison to the highest peak in the MS spectrum.
PEP		Posterior Error Probability of the identification. This value essentially operates as a p-value, where smaller is more significant.

MS/MS count		The number of sequencing events for this sequence, which matches the number of identifiers stored in the column MS/MS IDs.
MS/MS scan number		The RAW-file derived scan number of the MS/MS with the highest peptide identification score (the highest score is stored in the column 'Score').
MS/MS scan numbers		The scan numbers of all MS/MS spectra that are associated with this evidence.
MS3 scan numbers		The scan numbers of all MS3 spectra that are associated with this evidence.
Score		Andromeda score for the best associated MS/MS spectrum.
Delta score		Score difference to the second best identified peptide.
Combinatorics		Number of possible distributions of the modifications over the peptide sequence.
Ratio H/L		The ratio between two heavy and light label partners.
Ratio H/L normalized		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L shift		
Intensity		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Reverse		When marked with '+', this particular peptide was found to be part of a protein derived from the reversed part of the decoy database. These should be removed for further data analysis.
Potential contaminant		When marked with '+', this particular peptide was found to be part of a commonly occurring contaminant. These should be removed for further data analysis.
id		A unique (consecutive) identifier for each row in the evidence table, which is used to cross-link the information in this file with the information stored in the other files.
Protein group IDs		The identifier of the protein-group this redundant peptide sequence is associated with, which can be used to look up the extended protein information in the file 'proteinGroups.txt'. As a single peptide can be linked to multiple proteins (e.g. in the case of razor-proteins), multiple ids can be stored here separated by a semicolon. As a protein can be identified by multiple peptides, the same id can be found in different rows.
Peptide ID		The identifier of the non-redundant peptide sequence.
Mod. peptide ID		Identifier of the associated modification summary stored in the file 'modificationSpecificPeptides.txt'.
MS/MS IDs		Identifier(s) of the associated MS/MS summary(s) stored in the file 'msms.txt'.
Best MS/MS		Identifier(s) of the best MS/MS associated spectrum stored in the file 'msms.txt'.
Acetyl (K) site IDs		Identifier(s) of the modification summary stored in the file 'Acetyl (K)Sites.txt'.
Oxidation (M) site IDs		Identifier(s) of the modification summary stored in the file 'Oxidation (M)Sites.txt'.
Taxonomy IDs		Taxonomy identifiers.

# Peptides

The peptides table contains information on the identified peptides in the processed raw-files.

Name	Separator	Description
Sequence		The amino acid sequence of the identified peptide.
N-term cleavage window		Sequence window from -15 to 15 around the N-terminal cleavage site of this peptide.
C-term cleavage window		Sequence window from -15 to 15 around the C-terminal cleavage site of this peptide.
Amino acid before		The amino acid in the protein sequence before the peptide.
First amino acid		The amino acid in the first position of the peptide sequence.
Second amino acid		The amino acid in the first position of the peptide sequence.
Second last amino acid		The amino acid in the last position of the peptide sequence.
Last amino acid		The amino acid in the last position of the peptide sequence.
Amino acid after		The amino acid in the protein sequence after the peptide.
A Count		The number of instances of the 'A' amino acid contained within the sequence.
R Count		The number of instances of the 'R' amino acid contained within the sequence.
N Count		The number of instances of the 'N' amino acid contained within the sequence.
D Count		The number of instances of the 'D' amino acid contained within the sequence.
C Count		The number of instances of the 'C' amino acid contained within the sequence.
Q Count		The number of instances of the 'Q' amino acid contained within the sequence.
E Count		The number of instances of the 'E' amino acid contained within the sequence.
G Count		The number of instances of the 'G' amino acid contained within the sequence.
H Count		The number of instances of the 'H' amino acid contained within the sequence.
I Count		The number of instances of the 'I' amino acid contained within the sequence.
L Count		The number of instances of the 'L' amino acid contained within the sequence.
K Count		The number of instances of the 'K' amino acid contained within the sequence.
M Count		The number of instances of the 'M' amino acid contained within the sequence.
F Count		The number of instances of the 'F' amino acid contained within the sequence.
P Count		The number of instances of the 'P' amino acid contained within the sequence.
S Count		The number of instances of the 'S' amino acid contained within the sequence.
T Count		The number of instances of the 'T' amino acid contained within the sequence.
W Count		The number of instances of the 'W' amino acid contained within the sequence.
Y Count		The number of instances of the 'Y' amino acid contained within the sequence.
V Count		The number of instances of the 'V' amino acid contained within the sequence.
U Count		The number of instances of the 'U' amino acid contained within the sequence.
O Count		The number of instances of the 'O' amino acid contained within the sequence.
Length		The length of the sequence stored in the column "Sequence".
Missed cleavages		Number of missed enzymatic cleavages.
Mass		Monoisotopic mass of the peptide.
Proteins		Identifiers of proteins this peptide is associated with.
Leading razor protein		Identifier of the leading protein in the protein group which uses this peptide for quantification. (Either unique or razor.)
Start position		Position of the first amino acid of this peptide in the protein sequence. (one-based)
End position		Position of the last amino acid of this peptide in the protein sequence. (one-based)

Gene names		Names of genes this peptide is associated with.
Protein names		Names of proteins this peptide is associated with.
Unique (Groups)		When marked with '+', this particular peptide is unique to a single protein group in the proteinGroups file.
Unique (Proteins)		When marked with '+', this particular peptide is unique to a single protein sequence in the fasta file(s).
Charges		All charge states that have been observed.
PEP		Posterior Error Probability of the identification. This value essentially operates as a p-value, where smaller is more significant.
Score		Highest Andromeda score for the associated MS/MS spectra.
Identification type JSi09		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Identification type JSi10		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Identification type JSi11		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Identification type JSi12		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Fraction Average		
Fraction Std. Dev.		
Fraction 1		
Fraction 2		
Fraction 3		
Fraction 11		
Fraction 12		
Fraction 13		
Fraction 14		
Fraction 15		
Fraction 16		
Experiment JSi09		Number of evidence entries for this 'Experiment'.
Experiment JSi10		Number of evidence entries for this 'Experiment'.
Experiment JSi11		Number of evidence entries for this 'Experiment'.
Experiment JSi12		Number of evidence entries for this 'Experiment'.
Ratio H/L		The ratio between two heavy and light label partners.
Ratio H/L normalized		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L variability [%]		Coefficient of variability over all redundant quantifiable peptides. It is calculated as the standard deviation of the natural logarithm of ratios times 100.
Ratio H/L count		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type		
Ratio H/L JSi09		The ratio between two heavy and light label partners.
Ratio H/L normalized JSi09		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L variability [%] JSi09		Coefficient of variability over all redundant quantifiable peptides. It is calculated as the standard deviation of the natural logarithm of ratios times 100.
Ratio H/L count JSi09		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count JSi09		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type JSi09		
Ratio H/L JSi10		The ratio between two heavy and light label partners.
Ratio H/L normalized JSi10		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L variability [%] JSi10		Coefficient of variability over all redundant quantifiable peptides. It is calculated as the standard deviation of the natural logarithm of ratios times 100.
Ratio H/L count JSi10		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count JSi10		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type JSi10		
Ratio H/L JSi11		The ratio between two heavy and light label partners.
Ratio H/L normalized JSi11		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.

Ratio H/L variability [%] JSi11		Coefficient of variability over all redundant quantifiable peptides. It is calculated as the standard deviation of the natural logarithm of ratios times 100.
Ratio H/L count JSi11		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count JSi11		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type JSi11		
Ratio H/L JSi12		The ratio between two heavy and light label partners.
Ratio H/L normalized JSi12		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L variability [%] JSi12		Coefficient of variability over all redundant quantifiable peptides. It is calculated as the standard deviation of the natural logarithm of ratios times 100.
Ratio H/L count JSi12		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count JSi12		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type JSi12		
Intensity		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Intensity JSi09		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi09		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H JSi09		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Intensity JSi10		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi10		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H JSi10		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Intensity JSi11		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi11		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H JSi11		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Intensity JSi12		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi12		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H JSi12		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Reverse		When marked with '+', this particular peptide was found to be part of a protein derived from the reversed part of the decoy database. These should be removed for further data analysis.
Potential contaminant		When marked with '+', this particular peptide was found to be part of a commonly occurring contaminant. These should be removed for further data analysis.
id		A unique (consecutive) identifier for each row in the peptides table, which is used to cross-link the information in this table with the information stored in the other tables.
Protein group IDs		The identifiers of the protein groups this peptide was linked to, referenced against the proteinGroups table.
Mod. peptide IDs		Identifier(s) for peptide sequence(s), associated with the peptide, referenced against the corresponding modified peptides table.
Evidence IDs		Identifier(s) for analyzed peptide evidence associated with the protein group referenced against the evidence table.
MS/MS IDs		The identifiers of the MS/MS scans identifying this peptide, referenced against the msms table.
Best MS/MS		The identifier of the best (in terms of quality) MS/MS scan identifying this peptide, referenced against the msms table.



Acetyl (K) site IDs		Identifier(s) for site(s) associated with the protein group, which show(s) evidence of the modification, referenced against the appropriate modification site file.
Oxidation (M) site IDs		Identifier(s) for site(s) associated with the protein group, which show(s) evidence of the modification, referenced against the appropriate modification site file.
Taxonomy IDs		Taxonomy identifiers.
MS/MS Count		

# Modification-specific peptides

Name	Separator	Description
Sequence		The identified AA sequence of the peptide.
K Count		The number of instances of the 'K' AA contained within the sequence. The value for this can reliably be determined in the case of SILAC partners based on the distance between the partners. These counts are used to solidify the peptide identification process.
Modifications		Post-translational modifications contained within the sequence. When no modifications exist, this is set to 'unmodified'.
Mass		Charge corrected mass of the precursor ion.
Mass Fractional Part		The values after the decimal point (ie value - floor(value)).
Protein Groups		IDs of the protein groups to which this peptide belongs.
Proteins		The identifiers of the proteins this particular peptide is associated with.
Gene Names		Names of genes this peptide is associated with.
Protein Names		Names of proteins this peptide is associated with.
Unique (Groups)		When marked with '+', this particular peptide is unique to a single protein group in the proteinGroups file.
Unique (Proteins)		When marked with '+', this particular peptide is unique to a single protein sequence in the fasta file(s).
Acetyl (K)		Number of Acetyl (K) on this peptide.
Acetyl (Protein N-term)		Number of Acetyl (Protein N-term) on this peptide.
Oxidation (M)		Number of Oxidation (M) on this peptide.
Missed cleavages		Number of missed enzymatic cleavages.
Identification type JSi09		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Identification type JSi10		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Identification type JSi11		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Identification type JSi12		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Fraction Average		
Fraction Std. Dev.		
Fraction 1		
Fraction 2		
Fraction 3		
Fraction 11		
Fraction 12		
Fraction 13		
Fraction 14		
Fraction 15		
Fraction 16		
Experiment JSi09		Number of evidence entries for this 'Experiment'.
Experiment JSi10		Number of evidence entries for this 'Experiment'.
Experiment JSi11		Number of evidence entries for this 'Experiment'.
Experiment JSi12		Number of evidence entries for this 'Experiment'.
Retention time		Retention time in minutes averaged over the evidence entries belonging to this modification-specific peptide.
Calibrated retention time		Calibrated retention time averaged over the evidence entries belonging to this modification-specific peptide. Obviously this only makes sense if retention time recalibration has been performed which is the case when matching between run is selected.
Charges		All charge states that have been observed.
PEP		Posterior Error Probability of the identification. This value essentially operates as a p-value, where smaller is more significant.
MS/MS scan number		The RAW-file derived scan number of the MS/MS with the highest peptide identification score (the highest score is stored in the column 'Score').
Raw file		The name of the RAW-file the mass spectral data was derived from.
Score		Andromeda score for the best identified among the associated MS/MS spectra.
Delta score		Score difference to the second best identified peptide.

Reverse		When marked with '+', this particular peptide was found to be part of a protein derived from the reversed part of the decoy database. These should be removed for further data analysis.
Potential contaminant		When marked with '+', this particular peptide was found to be part of a commonly occurring contaminant. These should be removed for further data analysis.
Ratio H/L		The ratio between two heavy and light label partners.
Ratio H/L normalized		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L variability [%]		Coefficient of variability over all redundant quantifiable peptides. It is calculated as the standard deviation of the natural logarithm of ratios times 100.
Ratio H/L count		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type		
Ratio H/L JSi09		The ratio between two heavy and light label partners.
Ratio H/L normalized JSi09		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L variability [%] JSi09		Coefficient of variability over all redundant quantifiable peptides. It is calculated as the standard deviation of the natural logarithm of ratios times 100.
Ratio H/L count JSi09		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count JSi09		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type JSi09		
Ratio H/L JSi10		The ratio between two heavy and light label partners.
Ratio H/L normalized JSi10		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L variability [%] JSi10		Coefficient of variability over all redundant quantifiable peptides. It is calculated as the standard deviation of the natural logarithm of ratios times 100.
Ratio H/L count JSi10		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count JSi10		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type JSi10		
Ratio H/L JSi11		The ratio between two heavy and light label partners.
Ratio H/L normalized JSi11		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L variability [%] JSi11		Coefficient of variability over all redundant quantifiable peptides. It is calculated as the standard deviation of the natural logarithm of ratios times 100.
Ratio H/L count JSi11		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count JSi11		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type JSi11		
Ratio H/L JSi12		The ratio between two heavy and light label partners.
Ratio H/L normalized JSi12		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L variability [%] JSi12		Coefficient of variability over all redundant quantifiable peptides. It is calculated as the standard deviation of the natural logarithm of ratios times 100.
Ratio H/L count JSi12		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count JSi12		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type JSi12		
Intensity		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Intensity JSi09		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi09		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H JSi09		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.

Intensity JSi10		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi10		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H JSi10		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Intensity JSi11		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi11		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H JSi11		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Intensity JSi12		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi12		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H JSi12		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
id		A unique (consecutive) identifier for each row in the peptides table, which is used to cross-link the information in this table with the information stored in the other tables.
Protein group IDs		The identifiers of the protein groups this peptide was linked to, referenced against the proteinGroups table.
Peptide ID		Identifier of the associated peptide sequence summary, which can be found in the file 'peptides.txt'.
Evidence IDs		Identifier(s) for analyzed peptide evidence associated with the protein group referenced against the evidence table.
MS/MS IDs		The identifiers of the MS/MS scans identifying this peptide, referenced against the msms table.
Best MS/MS		The identifier of the best (in terms of quality) MS/MS scan identifying this peptide, referenced against the msms table.
Acetyl (K) site IDs		Identifier(s) for site(s) associated with this peptide, which show(s) evidence of the modification, referenced against the appropriate modification site file.
Oxidation (M) site IDs		Identifier(s) for site(s) associated with this peptide, which show(s) evidence of the modification, referenced against the appropriate modification site file.
MS/MS Count		
Taxonomy IDs		Taxonomy identifiers.

# Acetyl (K)Sites

Name	Separator	Description
Proteins		Identifiers of proteins this site is associated with.
Positions within proteins		For each protein identifier in the 'Proteins' column you find here the position of the site in the respective protein sequence. The index of the first amino acid in the sequence is 1.
Leading proteins		
Protein		Identifier of the protein this peptide is associated with.
Protein names		Names of proteins this peptide is associated with.
Gene names		Names of genes this peptide is associated with.
Fasta headers		Descriptions of proteins this peptide is associated with.
Localization prob		
Score diff		
PEP		The posterior error probability (PEP) of the best identified modified peptide containing this site.
Score		The Andromeda score of the best identified modified peptide containing this site.
Delta score		The Andromeda delta score of the best identified modified peptide containing this site.
Score for localization		The Andromeda score of the MS/MS spectrum used for calculating the localization score for this site.
Localization prob JSi09		
Score diff JSi09		
PEP JSi09		
Score JSi09		
Localization prob JSi10		
Score diff JSi10		
PEP JSi10		
Score JSi10		
Localization prob JSi11		
Score diff JSi11		
PEP JSi11		
Score JSi11		
Localization prob JSi12		
Score diff JSi12		
PEP JSi12		
Score JSi12		
Diagnostic peak		
Number of Acetyl (K)		Different numbers of Acetyl (K) on peptides that this site is involved in.
Amino acid		
Sequence window		
Modification window		
Peptide window coverage		
Acetyl (K) Probabilities		
Acetyl (K) Score diffs		
Position in peptide		
Charge		Charge state of the precursor ion.
Mass error [ppm]		Mass error of the recalibrated mass-over-charge value of the precursor ion in comparison to the predicted monoisotopic mass of the identified peptide sequence.
Identification type JSi09		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Identification type JSi10		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Identification type JSi11		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Identification type JSi12		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Ratio H/L		The ratio between two heavy and light label partners.
Ratio H/L 1		The ratio between two heavy and light label partners.
Ratio H/L 2		The ratio between two heavy and light label partners.
Ratio H/L 3		The ratio between two heavy and light label partners.

Ratio H/L normalized		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized___1		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized___2		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized___3		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L unmod. pep.		
Ratio H/L localized		
Ratio H/L variability [%]		Coefficient of variability over all redundant quantifiable peptides. It is calculated as the standard deviation of the natural logarithm of ratios times 100.
Ratio H/L count		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type		
Occupancy L		
Occupancy H		
Ratio H/L JSi09		The ratio between two heavy and light label partners.
Ratio H/L JSi09___1		The ratio between two heavy and light label partners.
Ratio H/L JSi09___2		The ratio between two heavy and light label partners.
Ratio H/L JSi09___3		The ratio between two heavy and light label partners.
Ratio H/L normalized JSi09		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi09___1		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi09___2		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi09___3		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L unmod. pep. JSi09		
Ratio H/L localized JSi09		
Ratio H/L variability [%] JSi09		
Ratio H/L count JSi09		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count JSi09		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type JSi09		
Occupancy L JSi09		
Occupancy H JSi09		
Ratio H/L JSi10		The ratio between two heavy and light label partners.
Ratio H/L JSi10___1		The ratio between two heavy and light label partners.
Ratio H/L JSi10___2		The ratio between two heavy and light label partners.
Ratio H/L JSi10___3		The ratio between two heavy and light label partners.
Ratio H/L normalized JSi10		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi10___1		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi10___2		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi10___3		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L unmod. pep. JSi10		
Ratio H/L localized JSi10		
Ratio H/L variability [%] JSi10		
Ratio H/L count JSi10		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count JSi10		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type JSi10		
Occupancy L JSi10		
Occupancy H JSi10		
Ratio H/L JSi11		The ratio between two heavy and light label partners.
Ratio H/L JSi11___1		The ratio between two heavy and light label partners.
Ratio H/L JSi11___2		The ratio between two heavy and light label partners.
Ratio H/L JSi11___3		The ratio between two heavy and light label partners.
Ratio H/L normalized JSi11		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.

Ratio H/L normalized JSi11___1		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi11___2		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi11___3		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L unmod. pep. JSi11		
Ratio H/L localized JSi11		
Ratio H/L variability [%] JSi11		
Ratio H/L count JSi11		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count JSi11		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type JSi11		
Occupancy L JSi11		
Occupancy H JSi11		
Ratio H/L JSi12		The ratio between two heavy and light label partners.
Ratio H/L JSi12___1		The ratio between two heavy and light label partners.
Ratio H/L JSi12___2		The ratio between two heavy and light label partners.
Ratio H/L JSi12___3		The ratio between two heavy and light label partners.
Ratio H/L normalized JSi12		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi12___1		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi12___2		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi12___3		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L unmod. pep. JSi12		
Ratio H/L localized JSi12		
Ratio H/L variability [%] JSi12		
Ratio H/L count JSi12		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count JSi12		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type JSi12		
Occupancy L JSi12		
Occupancy H JSi12		
Intensity		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Ratio mod/base L		
Ratio mod/base H		
Intensity JSi09		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi09		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H JSi09		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Ratio mod/base L JSi09		
Ratio mod/base H JSi09		
Intensity JSi10		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi10		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H JSi10		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Ratio mod/base L JSi10		
Ratio mod/base H JSi10		
Intensity JSi11		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi11		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.

Intensity H JSi11		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Ratio mod/base L JSi11		
Ratio mod/base H JSi11		
Intensity JSi12		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi12		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H JSi12		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Ratio mod/base L JSi12		
Ratio mod/base H JSi12		
Occupancy JSi09		
Occupancy ratioJSi09		
Occupancy error scale JSi09		
Occupancy JSi10		
Occupancy ratioJSi10		
Occupancy error scale JSi10		
Occupancy JSi11		
Occupancy ratioJSi11		
Occupancy error scale JSi11		
Occupancy JSi12		
Occupancy ratioJSi12		
Occupancy error scale JSi12		
Reverse		When marked with '+', this particular peptide was found to be part of a protein derived from the reversed part of the protein sequence database. These should be removed for further data analysis.
Potential contaminant		When marked with '+', this particular peptide was found to be part of a commonly occurring contaminant. These should be removed for further data analysis.
id		A unique (consecutive) identifier for each row in the site table, which is used to cross-link the information in this file with the information stored in the other files.
Protein group IDs		The identifier of the protein-group this peptide sequence is associated with, which can be used to look up the extended protein information in the file 'proteinGroups.txt'. As a single peptide can be linked to multiple proteins (e.g. in the case of razor-proteins), multiple id's can be stored here separated by a semicolon. As a protein can be identified by multiple peptides, the same id can be found in different rows.
Positions		The positions of the modifications in the protein amino acid sequence.
Position		The position of the modification in the protein amino acid sequence.
Peptide IDs		Identifier(s) of the associated peptide sequence(s) summary, which can be found in the file 'peptides.txt'.
Mod. peptide IDs		Identifier(s) of the associated peptide sequence(s) summary, which can be found in the file 'modificationSpecificPeptides.txt'.
Evidence IDs		Identifier(s) for analyzed peptide evidence associated with the protein group referenced against the evidence table.
MS/MS IDs		The identifiers of the MS/MS scans identifying this peptide, referenced against the msms table.
Best localization evidence ID		
Best localization MS/MS ID		
Best localization raw file		
Best localization scan number		
Best score evidence ID		
Best score MS/MS ID		
Best score raw file		
Best score scan number		
Best PEP evidence ID		
Best PEP MS/MS ID		
Best PEP raw file		
Best PEP scan number		



# Oxidation (M) Sites

Name	Separator	Description
Proteins		Identifiers of proteins this site is associated with.
Positions within proteins		For each protein identifier in the 'Proteins' column you find here the position of the site in the respective protein sequence. The index of the first amino acid in the sequence is 1.
Leading proteins		
Protein		Identifier of the protein this peptide is associated with.
Protein names		Names of proteins this peptide is associated with.
Gene names		Names of genes this peptide is associated with.
Fasta headers		Descriptions of proteins this peptide is associated with.
Localization prob		
Score diff		
PEP		The posterior error probability (PEP) of the best identified modified peptide containing this site.
Score		The Andromeda score of the best identified modified peptide containing this site.
Delta score		The Andromeda delta score of the best identified modified peptide containing this site.
Score for localization		The Andromeda score of the MS/MS spectrum used for calculating the localization score for this site.
Localization prob JSi09		
Score diff JSi09		
PEP JSi09		
Score JSi09		
Localization prob JSi10		
Score diff JSi10		
PEP JSi10		
Score JSi10		
Localization prob JSi11		
Score diff JSi11		
PEP JSi11		
Score JSi11		
Localization prob JSi12		
Score diff JSi12		
PEP JSi12		
Score JSi12		
Diagnostic peak		
Number of Oxidation (M)		Different numbers of Oxidation (M) on peptides that this site is involved in.
Amino acid		
Sequence window		
Modification window		
Peptide window coverage		
Oxidation (M) Probabilities		
Oxidation (M) Score diffs		
Position in peptide		
Charge		Charge state of the precursor ion.
Mass error [ppm]		Mass error of the recalibrated mass-over-charge value of the precursor ion in comparison to the predicted monoisotopic mass of the identified peptide sequence.
Identification type JSi09		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Identification type JSi10		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Identification type JSi11		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Identification type JSi12		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Ratio H/L		The ratio between two heavy and light label partners.
Ratio H/L 1		The ratio between two heavy and light label partners.
Ratio H/L 2		The ratio between two heavy and light label partners.
Ratio H/L 3		The ratio between two heavy and light label partners.

Ratio H/L normalized		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized___1		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized___2		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized___3		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L unmod. pep.		
Ratio H/L localized		
Ratio H/L variability [%]		Coefficient of variability over all redundant quantifiable peptides. It is calculated as the standard deviation of the natural logarithm of ratios times 100.
Ratio H/L count		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type		
Occupancy L		
Occupancy H		
Ratio H/L JSi09		The ratio between two heavy and light label partners.
Ratio H/L JSi09___1		The ratio between two heavy and light label partners.
Ratio H/L JSi09___2		The ratio between two heavy and light label partners.
Ratio H/L JSi09___3		The ratio between two heavy and light label partners.
Ratio H/L normalized JSi09		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi09___1		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi09___2		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi09___3		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L unmod. pep. JSi09		
Ratio H/L localized JSi09		
Ratio H/L variability [%] JSi09		
Ratio H/L count JSi09		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count JSi09		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type JSi09		
Occupancy L JSi09		
Occupancy H JSi09		
Ratio H/L JSi10		The ratio between two heavy and light label partners.
Ratio H/L JSi10___1		The ratio between two heavy and light label partners.
Ratio H/L JSi10___2		The ratio between two heavy and light label partners.
Ratio H/L JSi10___3		The ratio between two heavy and light label partners.
Ratio H/L normalized JSi10		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi10___1		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi10___2		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi10___3		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L unmod. pep. JSi10		
Ratio H/L localized JSi10		
Ratio H/L variability [%] JSi10		
Ratio H/L count JSi10		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count JSi10		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type JSi10		
Occupancy L JSi10		
Occupancy H JSi10		
Ratio H/L JSi11		The ratio between two heavy and light label partners.
Ratio H/L JSi11___1		The ratio between two heavy and light label partners.
Ratio H/L JSi11___2		The ratio between two heavy and light label partners.
Ratio H/L JSi11___3		The ratio between two heavy and light label partners.
Ratio H/L normalized JSi11		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.

Ratio H/L normalized JSi11___1		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi11___2		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi11___3		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L unmod. pep. JSi11		
Ratio H/L localized JSi11		
Ratio H/L variability [%] JSi11		
Ratio H/L count JSi11		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count JSi11		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type JSi11		
Occupancy L JSi11		
Occupancy H JSi11		
Ratio H/L JSi12		The ratio between two heavy and light label partners.
Ratio H/L JSi12___1		The ratio between two heavy and light label partners.
Ratio H/L JSi12___2		The ratio between two heavy and light label partners.
Ratio H/L JSi12___3		The ratio between two heavy and light label partners.
Ratio H/L normalized JSi12		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi12___1		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi12___2		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi12___3		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L unmod. pep. JSi12		
Ratio H/L localized JSi12		
Ratio H/L variability [%] JSi12		
Ratio H/L count JSi12		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count JSi12		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type JSi12		
Occupancy L JSi12		
Occupancy H JSi12		
Intensity		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Ratio mod/base L		
Ratio mod/base H		
Intensity JSi09		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi09		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H JSi09		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Ratio mod/base L JSi09		
Ratio mod/base H JSi09		
Intensity JSi10		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi10		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H JSi10		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Ratio mod/base L JSi10		
Ratio mod/base H JSi10		
Intensity JSi11		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi11		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.

Intensity H JSi11		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Ratio mod/base L JSi11		
Ratio mod/base H JSi11		
Intensity JSi12		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi12		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H JSi12		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Ratio mod/base L JSi12		
Ratio mod/base H JSi12		
Reverse		When marked with '+', this particular peptide was found to be part of a protein derived from the reversed part of the protein sequence database. These should be removed for further data analysis.
Potential contaminant		When marked with '+', this particular peptide was found to be part of a commonly occurring contaminant. These should be removed for further data analysis.
id		A unique (consecutive) identifier for each row in the site table, which is used to cross-link the information in this file with the information stored in the other files.
Protein group IDs		The identifier of the protein-group this peptide sequence is associated with, which can be used to look up the extended protein information in the file 'proteinGroups.txt'. As a single peptide can be linked to multiple proteins (e.g. in the case of razor-proteins), multiple id's can be stored here separated by a semicolon. As a protein can be identified by multiple peptides, the same id can be found in different rows.
Positions		The positions of the modifications in the protein amino acid sequence.
Position		The position of the modification in the protein amino acid sequence.
Peptide IDs		Identifier(s) of the associated peptide sequence(s) summary, which can be found in the file 'peptides.txt'.
Mod. peptide IDs		Identifier(s) of the associated peptide sequence(s) summary, which can be found in the file 'modificationSpecificPeptides.txt'.
Evidence IDs		Identifier(s) for analyzed peptide evidence associated with the protein group referenced against the evidence table.
MS/MS IDs		The identifiers of the MS/MS scans identifying this peptide, referenced against the msms table.
Best localization evidence ID		
Best localization MS/MS ID		
Best localization raw file		
Best localization scan number		
Best score evidence ID		
Best score MS/MS ID		
Best score raw file		
Best score scan number		
Best PEP evidence ID		
Best PEP MS/MS ID		
Best PEP raw file		
Best PEP scan number		

# Protein groups

The Protein Groups table contains information about the identified proteins in the processed raw-files. Each row contains refers to a group of proteins that could be reconstructed from a set of peptides.

Name	Separator	Description
Protein IDs		Identifiers of proteins contained in the protein group. They are sorted by number of identified peptides in descending order.
Majority protein IDs		These are the IDs of those proteins that have at least half of the peptides that the leading protein has.
Peptide counts (all)		Number of peptides associated with each protein in protein group, occurring in the order as the protein IDs occur in the 'Protein IDs' column. Here distinct peptide sequences are counted. Modified forms or different charges are counted as one peptide.
Peptide counts (razor+unique)		Number of peptides associated with each protein in protein group, occurring in the order as the protein IDs occur in the 'Protein IDs' column. Here distinct peptide sequences are counted. Modified forms or different charges are counted as one peptide.
Peptide counts (unique)		Number of peptides associated with each protein in protein group, occurring in the order as the protein IDs occur in the 'Protein IDs' column. Here distinct peptide sequences are counted. Modified forms or different charges are counted as one peptide.
Protein names		Names of proteins contained in the group.
Gene names		Names of the genes associated to the proteins contained in the group.
Fasta headers		Fasta headers(s) of protein(s) contained in the group.
Number of proteins		Number of proteins contained within the group. This corresponds to the number of entries in the column 'Protein IDs'.
Peptides		The total number of peptide sequences associated with the protein group (i.e. for all the proteins in the group).
Razor + unique peptides		The total number of razor + unique peptides associated with the protein group (i.e. these peptides are shared with another protein group).
Unique peptides		The total number of unique peptides associated with the protein group (i.e. these peptides are not shared with another protein group).
Peptides JSi09		Number of peptides (distinct peptide sequences) in experiment JSi09
Peptides JSi10		Number of peptides (distinct peptide sequences) in experiment JSi10
Peptides JSi11		Number of peptides (distinct peptide sequences) in experiment JSi11
Peptides JSi12		Number of peptides (distinct peptide sequences) in experiment JSi12
Razor + unique peptides JSi09		Number of razor + unique peptides (distinct peptide sequences) in experiment JSi09
Razor + unique peptides JSi10		Number of razor + unique peptides (distinct peptide sequences) in experiment JSi10
Razor + unique peptides JSi11		Number of razor + unique peptides (distinct peptide sequences) in experiment JSi11
Razor + unique peptides JSi12		Number of razor + unique peptides (distinct peptide sequences) in experiment JSi12
Unique peptides JSi09		Number of unique peptides (distinct peptide sequences) in experiment JSi09
Unique peptides JSi10		Number of unique peptides (distinct peptide sequences) in experiment JSi10
Unique peptides JSi11		Number of unique peptides (distinct peptide sequences) in experiment JSi11
Unique peptides JSi12		Number of unique peptides (distinct peptide sequences) in experiment JSi12
Sequence coverage [%]		Percentage of the sequence that is covered by the identified peptides of the first protein sequence contained in the group.
Unique + razor sequence coverage [%]		Percentage of the sequence that is covered by the identified unique and razor peptides of the best protein sequence contained in the group.
Unique sequence coverage [%]		Percentage of the sequence that is covered by the identified unique peptides of the best protein sequence contained in the group.
Mol. weight [kDa]		Molecular weight of the leading protein sequence contained in the protein group.

Sequence length		The length of the leading protein sequence contained in the group.
Sequence lengths		The length of all sequences of the proteins contained in the group.
Fraction average		
Fraction 1		
Fraction 2		
Fraction 3		
Fraction 11		
Fraction 12		
Fraction 13		
Fraction 14		
Fraction 15		
Fraction 16		
Q-value		This is the ratio of reverse to forward protein groups.
Score		Protein score which is derived from peptide posterior error probabilities.
Identification type JSi09		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Identification type JSi10		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Identification type JSi11		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Identification type JSi12		Indicates whether this experiment was identified by MS/MS or only by matching between runs.
Ratio H/L		The ratio between two heavy and light label partners.
Ratio H/L normalized		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L variability [%]		Coefficient of variability over all redundant quantifiable peptides. It is calculated as the standard deviation of the natural logarithm of ratios times 100.
Ratio H/L count		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type		
Ratio H/L JSi09		The ratio between two heavy and light label partners.
Ratio H/L JSi10		The ratio between two heavy and light label partners.
Ratio H/L JSi11		The ratio between two heavy and light label partners.
Ratio H/L JSi12		The ratio between two heavy and light label partners.
Ratio H/L normalized JSi09		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi10		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi11		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L normalized JSi12		Normalized ratio between two medium and light label partners. The median of ratio sub-populations was shifted to 1.
Ratio H/L variability [%] JSi09		Coefficient of variability over all redundant quantifiable peptides. It is calculated as the standard deviation of the natural logarithm of ratios times 100.
Ratio H/L variability [%] JSi10		Coefficient of variability over all redundant quantifiable peptides. It is calculated as the standard deviation of the natural logarithm of ratios times 100.
Ratio H/L variability [%] JSi11		Coefficient of variability over all redundant quantifiable peptides. It is calculated as the standard deviation of the natural logarithm of ratios times 100.
Ratio H/L variability [%] JSi12		Coefficient of variability over all redundant quantifiable peptides. It is calculated as the standard deviation of the natural logarithm of ratios times 100.
Ratio H/L count JSi09		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L count JSi10		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L count JSi11		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L count JSi12		Number of redundant peptides (MS1 features) used for quantitation.
Ratio H/L iso-count JSi09		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L iso-count JSi10		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L iso-count JSi11		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.

Ratio H/L iso-count JSi12		Number of redundant peptides (MS1 features) used for quantitation that are quantified with the re-quantify method.
Ratio H/L type JSi09		
Ratio H/L type JSi10		
Ratio H/L type JSi11		
Ratio H/L type JSi12		
Sequence coverage JSi09 [%]		Percentage of the sequence that is covered by the identified peptides in this sample of the longest protein sequence contained within the group.
Sequence coverage JSi10 [%]		Percentage of the sequence that is covered by the identified peptides in this sample of the longest protein sequence contained within the group.
Sequence coverage JSi11 [%]		Percentage of the sequence that is covered by the identified peptides in this sample of the longest protein sequence contained within the group.
Sequence coverage JSi12 [%]		Percentage of the sequence that is covered by the identified peptides in this sample of the longest protein sequence contained within the group.
Intensity		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Intensity JSi09		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi09		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H JSi09		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Intensity JSi10		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi10		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H JSi10		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Intensity JSi11		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi11		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H JSi11		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Intensity JSi12		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L JSi12		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H JSi12		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Peptide sequences		Peptide sequences belonging to this protein group.
Only identified by site		When marked with '+', this particular protein group was identified only by a modification site.
Reverse		When marked with '+', this particular protein group contains no protein, made up of at least 50% of the peptides of the leading protein, with a peptide derived from the reversed part of the decoy database. These should be removed for further data analysis. The 50% rule is in place to prevent spurious protein hits to erroneously flag the protein group as reverse.
Potential contaminant		When marked with '+', this particular protein group was found to be a commonly occurring contaminant. It is recommended to remove them for further data analysis.
id		A unique (consecutive) identifier for each row in the proteinGroups table, which is used to cross-link the information in this file with the information stored in the other files.
Peptide IDs		Identifiers of the associated peptides, which correspond to rows in the peptides table.
Peptide is razor		Indicates for each peptide ID if it is a razor or group unique peptide (true) or a non unique non razor peptide (false).
Mod. peptide IDs		
Evidence IDs		

MS/MS IDs		
Best MS/MS		The identifier of the best (in terms of quality) MS/MS scans identifying the peptides of this protein, referenced against the msms table.
Acetyl (K) site IDs		Identifier(s) for site(s) associated with the protein group, which show(s) evidence of the modification, referenced against the appropriate modification site file.
Oxidation (M) site IDs		Identifier(s) for site(s) associated with the protein group, which show(s) evidence of the modification, referenced against the appropriate modification site file.
Acetyl (K) site positions		Positions of the sites in the leading protein of this group.
Oxidation (M) site positions		Positions of the sites in the leading protein of this group.
Taxonomy IDs		Taxonomy identifiers.



# All peptides

Name	Separator	Description
Raw file		Name of the raw file the spectral data was extracted from.
Type		The type of detection for the peptide. MULTI – A labeling multiplet was detected. ISO – An isotope pattern was detected.
Charge		The charge state of the peptide.
m/z		The mass divided by the charge of the charged peptide.
Mass		The mass of the neutral peptide ((m/z-proton) * charge).
Uncalibrated m/z		m/z before re-calibrations have been applied.
Number of data points		The number of data points (peak centroids) collected for this peptide feature.
Number of scans		The number of MS scans that the 3d peaks of this peptide feature are overlapping with.
Number of isotopic peaks		The number of isotopic peaks contained in this peptide feature.
PIF		Short for Parent Ion Fraction; indicates the fraction the target peak makes up of the total intensity in the inclusion window.
Mass fractional part		The values after the radix point (ie value - floor(value)).
Mass deficit		Empirically derived deviation measure to the next nearest integer scaled to center around 0. Can be used to visually detect contaminants in a plot setting Mass against this value.  $m*a+b - \text{round}(m*a+b)$ m: the peptide mass a: 0.99954 b: -0.04
Mass precision [ppm]		The precision of the mass detection of the peptide in parts-per-million.
Max intensity m/z 0		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Max intensity m/z 1		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Retention time		The retention time of the peak detected for the peptide measured in minutes.
Retention length		The total retention time width of the peak (last time point – first time point) in seconds.
Retention length (FWHM)		The full width at half maximum value retention time width of the peak in seconds.
Min scan number		The first scan number at which the peak was encountered.
Max scan number		The last scan number at which the peak was encountered.
Identified		When marked with '+' this particular MS/MS scan was identified as a peptide; when marked with '-' no identification was made.
Reverse		When marked with '+' this particular MS/MS scan was assigned to a decoy sequence.
MS/MS IDs		Unique identifier linking this identification to the MS/MS scans.
Sequence		The identified AA sequence of the peptide.
Length		The length of the sequence stored in the column "Sequence".
Modifications		Post-translational modifications contained within the sequence. When no modifications exist, this is set to 'unmodified'.  Note: This column only set when this MS/MS spectrum has been identified.
Modified sequence		Sequence representation of the peptide including location(s) of modified AAs.  Note: This column only set when this MS/MS spectrum has been identified.
Proteins		Identifiers of proteins this peptide is associated with.  Note: This column only set when this MS/MS spectrum has been identified.
Score		The score of the identification (higher is better).
PEP		The posterior error probability of the identification (smaller is better).

Lys Count		The number of instances of Lys contained within the sequence. The value for this can reliably be determined in the case of label partners, based on the distance between the partners. These counts are used to solidify the peptide identification process.
Ratio H/L		The ratio between two heavy and light multiplet members.
Ratio H/L normalized		Normalized ratio between two heavy and light multiplet members. The median of the total ratio population was shifted to 1.
Intensity		Summed up eXtracted Ion Current (XIC) of all isotopic clusters associated with the identified AA sequence. In case of a labeled experiment this is the total intensity of all the isotopic patterns in the label cluster.
Intensity L		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the light label partner.
Intensity H		Summed up eXtracted Ion Current (XIC) of the isotopic cluster belonging to the heavy label partner.
Intensities L		Elution profile of the light peptide.
Intensities H		Elution profile of the heavy peptide.
Isotope pattern L		Isotope pattern of the light peptide.
Isotope pattern H		Isotope pattern of the heavy peptide.
MS/MS count		The number of MS/MS spectra recorded for the peptide.
MSMS scan numbers		The scan numbers where the MS/MS spectra were recorded.
MSMS label states		The label partner detected for the peptide. The value 0 is always the light partner. In the case of double label labeling 1 is the heavy partner. In the case of triple label labeling 1 is the medium and 2 the heavy partner.
MSMS isotope indices		Indices of the isotopic peaks that the MS/MS spectra reside on. A value of 0 corresponds to the monoisotopic peak.

# MS scans

The msScans table contains information about the full scans, which can be used to verify data quality and generated useful statistics about the interaction between the samples and LC.

Name	Separator	Description
Raw file		The name of the RAW-file the mass spectral data originates from.
Scan number		The scan number (defined in the raw-file) at which the full scan was made.
Scan index		The consecutive index of this full scan.
Retention time		The retention time at which the full scan was made.
Cycle time		The total time (full scan including the tandem MS scans) this full scan has taken up.
Ion injection time		The total injection time that was required to capture the specified amount of ions. This value is limited by a maximum, which can be used to determine whether the time has maxed out (indicative of a bad acquisition).
Base peak intensity		The intensity of the most intense ion in the spectrum.
Total ion current		The total intensity acquired in the full scan.
MS/MS count		The number of tandem MS scans that were made based on this full scan (e.g. a top 10 method selects the top 10 most intense ions in the scan and fragments those).
Mass calibration		The applied mass correction in Th to the full scan.
Fraction		The fraction measured with this full scan.
Experiment		
Peak length		The average time between the start and the end of the peaks detected in the full scan.
Isotope pattern length		The average time between the start and the end of the isotope patterns detected in the full scan.
Multiplet length		The average time between the start and the end of the isotope patterns of the labeling multiplets detected in the full scan.
Peaks / s		The average number of peaks detected per second of chromatography.
Single peaks / s		The average number of single peaks detected per second of chromatography.
Isotope patterns / s		The average number of isotope patterns detected per second of chromatography.
Single isotope patterns / s		The average number of single isotope patterns detected per second of chromatography.
Multiplets / s		The average number of labeling multiplets detected per second of chromatography.
Identified multiplets / s		The percentage of labeling multiplets actually identified.
Multiplet identification rate [%]		The percentage of the detected labeling multiplets that were identified.
MS/MS / s		The average number of MS/MS events per second of chromatography.
Identified MS/MS / s		The average number of identified MS/MS events per second of chromatography.
MS/MS identification rate [%]		The percentage of tandem MS scans that were identified.
Intens Comp Factor		Taken from the Thermo RAW file.
CTCD Comp		Taken from the Thermo RAW file.
RawOvFtT		For Thermo Fisher only. TIC estimation done with the orbitrap cell.
AGC Fill		Taken from the Thermo RAW file.

# MZ range

Name	Separator	Description
Raw file		The name of the RAW-file the mass spectral data was derived from.
m/z		The mass-over-charge value.
Peaks / Da		The average number of peaks detected per Dalton.
Single peaks / Da		The average number of single peaks detected per Dalton.
Isotope patterns / Da		The average number of isotope patterns detected per Dalton.
Single isotope patterns / Da		The average number of single isotope patterns detected per Dalton.
SILAC pairs / Da		The average number of SILAC pairs detected per Dalton.
Identified SILAC pairs / Da		The percentage of SILAC pairs actually identified.
SILAC identification rate [%]		The percentage of the detected SILAC pairs that were identified.
MS/MS / Da		The average number of MS/MS events per Dalton.
Identified MS/MS / Da		The average number of identified MS/MS events per Dalton.
Identification rate [%]		The percentage of tandem MS scans that were identified.

# MS/MS scans

Name	Separator	Description
Raw file		Name of the RAW file the spectral MS/MS data was extracted from.
Scan number		RAW file derived scan number for the MS/MS spectrum.
Retention time		Time point along the elution profile at which the MS/MS data was recorded.
Ion injection time		The ion inject time for the MS/MS scan. This can be used to determine if this time equals to the maximum ion inject time, general indicative of a lower quality spectrum.
Total ion current		The total ion current of the MS/MS scan. For Thermo data this value is calculated by summing all the intensity values found in the mass spectral data, which is different from the Xcalibur reported TIC (Xcalibur TIC is about 25% of the value reported here).
Collision energy		The collision energy used for the fragmentation that resulted in this MS/MS scan.
Summations		For time of flight instruments only.
Base peak intensity		The intensity of the most intense ion in the spectrum.
Elapsed time		The time the MS/MS scan took to complete.
Identified		When marked with '+' this particular MS/MS scan was identified as a peptide; when marked with '-' no identification was made.
Matched		When marked with '+' this particular MS/MS scan was retrieved by matching between runs.
Reverse		When marked with '+' this particular MS/MS scan was assigned to a decoy sequence.
MS/MS IDs		Unique identifier linking this identification to the MS/MS scans.
Sequence		The identified AA sequence of the peptide.
Length		The length of the sequence stored in the column "Sequence".
Filtered peaks		Number of peaks after the 'top X per 100 Da' filtering.
m/z		Recalibrated m/z of the precursor ion.
Mass		Charge corrected mass of the precursor ion.
Charge		Charge state of the precursor ion.
MS2 m/z		m/z value used for acquisition.
Type		The type of precursor ion as identified by MaxQuant. ISO – isotopic cluster. PEAK – single peak. MULTI – labeling cluster.
Fragmentation		The type of fragmentation used to create the MS/MS spectrum. CID – Collision Induced Dissociation. HCD – High energy Collision induced Dissociation. ETD – Electron Transfer Dissociation.
Mass analyzer		The mass analyzer used to record the MS/MS spectrum. ITMS – Ion trap. FTMS – Fourier transform ICR or orbitrap cell. TOF – Time of flight.
Parent intensity fraction		The percentage the parent ion intensity makes up of the total intensity in the selection window.
Fraction of total spectrum		The percentage the parent ion intensity makes up of the total intensity of the whole MS spectrum.
Base peak fraction		The percentage the parent ion intensity in comparison to the highest peak in the MS spectrum.
Precursor full scan number		The full scan number where the precursor ion was selected for fragmentation.
Precursor intensity		The intensity of the precursor ion at the scan number it was selected.
Precursor apex fraction		The fraction the intensity of the precursor ion makes up of the peak (apex) intensity.
Precursor apex offset		How many full scans the precursor ion is offset from the peak (apex) position.
Precursor apex offset time		How much time the precursor ion is offset from the peak (apex) position.
Scan event number		This number indicates which MS/MS scan this one is in the consecutive order of the MS/MS scans that are acquired after an MS scan.
Modifications		Post-translational modifications contained within the sequence. When no modifications exist, this is set to 'unmodified'.  Note: This column only set when this MS/MS spectrum has been identified.

Modified sequence		Sequence representation of the peptide including location(s) of modified AAs.  Note: This column only set when this MS/MS spectrum has been identified.
Proteins		Identifiers of proteins this peptide is associated with.  Note: This column only set when this MS/MS spectrum has been identified.
Score		The score of the identification (higher is better).
PEP		The posterior error probability of the identification (smaller is better).
Fraction		The identifier of the fraction the sample was taken from.
Experiment		
MS3 scan numbers		The scan numbers of all MS3 spectra that are associated with this MS/MS scan.
Reporter PIF		
Reporter fraction		
Intens Comp Factor		Taken from the Thermo RAW file.
CTCD Comp		Taken from the Thermo RAW file.
RawOvFtT		For Thermo Fisher only. TIC estimation done with the orbitrap cell.
AGC Fill		Taken from the Thermo RAW file.
Scan index		Consecutive index of the MS/MS spectrum.
MS scan index		Consecutive index of the MS spectrum prior to this MS/MS spectrum.
MS scan number		Scan number of the MS spectrum prior to this MS/MS spectrum.

# MS/MS

Name	Separator	Description
Raw file		The name of the RAW file the mass spectral data was read from.
Scan number		The RAW-file derived scan number of the MS/MS spectrum.
Scan index		The consecutive index of the MS/MS spectrum.
Sequence		The identified AA sequence of the peptide.
Length		The length of the sequence stored in the column "Sequence".
Missed cleavages		Number of missed enzymatic cleavages.
Modifications		Post-translational modifications contained within the identified peptide sequence.
Modified sequence		Sequence representation including the post-translational modifications (abbreviation of the modification in brackets before the modified AA). The sequence is always surrounded by underscore characters ('_').
Acetyl (K) Probabilities		Sequence representation of the peptide including PTM positioning probabilities ([0..1], where 1 is best match) for 'Acetyl (K)'.
Oxidation (M) Probabilities		Sequence representation of the peptide including PTM positioning probabilities ([0..1], where 1 is best match) for 'Oxidation (M)'.
Acetyl (K) Score diffs		
Oxidation (M) Score diffs		
Acetyl (K)		
Acetyl (Protein N-term)		
Oxidation (M)		
Proteins		The identifiers of the proteins the identified peptide is associated with.
Gene Names		Names of genes the identified peptide is associated with.
Protein Names		Descriptions of the proteins the identified peptide is associated with.
Charge		The charge state of the precursor ion.
Fragmentation		The type of fragmentation used to create the MS/MS spectrum. CID – Collision Induced Dissociation. HCD – High energy Collision induced Dissociation. ETD – Electron Transfer Dissociation.
Mass analyzer		The mass analyzer used to record the MS/MS spectrum. ITMS – Ion trap. FTMS – Fourier transform ICR or orbitrap cell. TOF – Time of flight.
Type		The type of precursor ion as identified by MaxQuant. ISO – isotopic cluster. PEAK – single peak. MULTI – labeling cluster.
Scan event number		
Isotope index		
m/z		The mass-over-charge of the precursor ion.
Mass		The charge corrected mass of the precursor ion.
Mass error [ppm]		Mass error of the recalibrated mass-over-charge value of the precursor ion in comparison to the predicted monoisotopic mass of the identified peptide sequence expressed in parts per million.
Mass error [Da]		Mass error of the recalibrated mass-over-charge value of the precursor ion in comparison to the predicted monoisotopic mass of the identified peptide sequence expressed in atomic mass units.
Simple mass error [ppm]		
Retention time		The uncalibrated retention time in minutes where the MS/MS spectrum has been acquired.
PEP		Posterior Error Probability of the identification. This value essentially operates as a p-value, where smaller is more significant.
Score		Andromeda score for the best associated MS/MS spectrum.
Delta score		Score difference to the second best identified peptide with a different amino acid sequence.
Score diff		Score difference to the second best positioning of modifications identified peptide with the same amino acid sequence.
Localization prob		
Combinatorics		Number of possible distributions of the modifications over the peptide sequence.

Labeling state		Labeling state of the precursor isotope pattern used to identify the peptide.
PIF		Short for Parent Ion Fraction; indicates the fraction the target peak makes up of the total intensity in the inclusion window.
Fraction of total spectrum		The percentage the parent ion intensity makes up of the total intensity of the whole spectrum.
Base peak fraction		The percentage the parent ion intensity in comparison to the highest peak in the MS spectrum.
Precursor full scan number		The full scan number where the precursor ion was selected for fragmentation.
Precursor Intensity		The intensity of the precursor ion at the scan number it was selected.
Precursor apex fraction		The fraction the intensity of the precursor ion makes up of the peak (apex) intensity.
Precursor apex offset		How many full scans the precursor ion is offset from the peak (apex) position.
Precursor apex offset time		How much time the precursor ion is offset from the peak (apex) position.
Diagnostic peak Acetyl (K) K		
Matches		The species of the peaks in the fragmentation spectrum after TopN filtering.
Intensities		The intensities of the peaks in the fragmentation spectrum after TopN filtering.
Mass deviations [Da]		The mass deviation of each peak in the fragmentation spectrum in absolute mass units.
Mass deviations [ppm]		The mass deviation of each peak in the fragmentation spectrum in parts per million.
Masses		The masses-over-charge of the peaks in the fragmentation spectrum.
Number of matches		The number of peaks matching to the predicted fragmentation spectrum.
Intensity coverage		The fraction of intensity in the MS/MS spectrum that is annotated.
Peak coverage		The fraction of peaks in the MS/MS spectrum that are annotated.
Neutral loss level		How many neutral losses were applied to each fragment in the Andromeda scoring.
ETD identification type		For ETD spectra several different combinations of ion series are scored. Here the highest scoring combination is indicated
Reverse		When marked with '+', this particular peptide was found to be part of a protein derived from the reversed part of the decoy database. These should be removed for further data analysis.
All scores		
All sequences		
All modified sequences		
MS3 scan numbers		The RAW-file derived scan number of the MS/MS spectrum.
Reporter PIF		
Reporter fraction		
id		A unique (consecutive) identifier for each row in the msms table, which is used to cross-link the information in this file with the information stored in the other files.
Protein group IDs		The identifier of the protein-group this redundant peptide sequence is associated with, which can be used to look up the extended protein information in the file 'proteinGroups.txt'. As a single peptide can be linked to multiple proteins (e.g. in the case of razor-proteins), multiple id's can be stored here separated by a semicolon. As a protein can be identified by multiple peptides, the same id can be found in different rows.
Peptide ID		The identifier of the non-redundant peptide sequence.
Mod. peptide ID		Identifier of the associated modification summary stored in the file 'modificationSpecificPeptides.txt'.
Evidence ID		Identifier of the associated evidence stored in the file 'evidence.txt'.
Acetyl (K) site IDs		Identifier of the associated entry stored in the file 'Acetyl (K)Sites.txt'.
Oxidation (M) site IDs		Identifier of the associated entry stored in the file 'Oxidation (M)Sites.txt'.